

The Hamburg/SAO Survey for Emission-Line Galaxies

V. The Fifth List of 161 Galaxies

A.V. Ugryumov^{1,11}, D. Engels², A.Y. Kniazev^{1,11}, R.F. Green³, Y.I. Izotov^{4,9}, U. Hopp^{5,10},
S.A. Pustilnik^{1,11}, A.G. Pramsky^{1,11}, T.F. Kniazeva¹, N. Brosch⁶, H.-J. Hagen², V.A. Lipovetsky^{1,*},
J. Masegosa⁷, I. Márquez⁷, and J.-M. Martín⁸

¹ Special Astrophysical Observatory, Nizhnij Arkhyz, Karachai-Circassia, 369167, Russia

² Hamburger Sternwarte, Gojenbergsweg 112, D-21029 Hamburg, Germany

³ National Optical Astronomical Observatories, Tucson, USA

⁴ Main Astronomical Observatory, Golosiiv, Kyiv-22, 03680, Ukraine

⁵ Universitätssternwarte München, Scheiner Str. 1, D-81679 München, Germany

⁶ Wise Observatory, Tel-Aviv University, Tel-Aviv 69978, Israel

⁷ Instituto de Astrofísica de Andalucía, CSIC, Aptdo. 3004, 18080, Granada, Spain

⁸ Département de Radioastronomie ARPEGES, Observatoire de Paris, F-92195 Meudon Cedex, France

⁹ Visiting astronomer at Kitt Peak National Observatory, USA

¹⁰ Visiting astronomer at Calar Alto Observatory, Spain

¹¹ Isaac Newton Institute of Chile, SAO Branch

Received ; Accepted

Abstract. We present the fifth list with results** of the Hamburg/SAO Survey for Emission-Line Galaxies (HSS therein, SAO – Special Astrophysical Observatory, Russia). The list is a result of follow-up spectroscopy conducted with the 2.2m CAHA and 4m Kitt Peak telescopes in 1999. The data of this snap-shot spectroscopy survey confirmed 166 emission-line objects out of 209 observed candidates and allowed their quantitative spectral classification and redshift determination. We could classify 98 emission-line objects as BCG/HII galaxies or probable BCGs, 5 – as QSOs, 3 – as Seyfert galaxies, 2 – as super-associations in subluminescent spiral galaxies. 25 low-excitation objects were classified as starburst nuclei (SBN), 24 as dwarf amorphous nuclei starburst galaxies (DANS) and 3 as LINERs. Due to low signal-to-noise ratio we could not classify 6 ELGs. Furthermore, for another 4 galaxies we did not detect any significant emission lines. For 131 emission-line galaxies, the redshifts and/or line intensities are determined for the first time. For the remaining 30 previously known ELGs we give either improved data of the line intensities or some independent measurements.

Key words. surveys – galaxies: fundamental parameters – galaxies: distances and redshifts – galaxies: starburst – galaxies: compact – quasars: redshifts

1. Introduction

The problem of creating large, homogeneous and deep samples of actively star-forming low-mass galaxies is very important for several applications in studies of galaxy evolution and spatial distribution. Several earlier projects, like the Second Byurakan Survey (SBS) (Markarian et al. 1983, Stepanian 1994), the University of Michigan

(UM) survey (e.g., Salzer et al. 1989), and the Case survey (Pesch et al. 1995, Salzer et al. 1995, Ugryumov et al. 1998), as well as some others, identified on objective prism plates many hundreds of emission-line galaxies. The Hamburg/SAO survey (HSS) is intended to create a new very large homogeneous sample of such galaxies in the region of the Northern sky with an area of some 1700 square degrees. The basic outline of the HSS and first results are described in Paper I (Ugryumov et al. 1999), while additional results from follow-up spectroscopy are given in papers II, III and IV (Pustilnik et al. 1999, Hopp et al. 2000, Kniazev et al. 2001a). In this paper we present the results of follow-up spectroscopy of another 209 objects selected on the Hamburg Quasar Survey (HQS) prism spectral plates as ELG candidates.

Send offprint requests to: A.Ugryumov e-mail: and@sao.ru

* Deceased 1996 September 22.

** Tables 3 to 7 are only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/Abstract.html>. Figures A1 to A17 will be made available only in the electronic version of the journal.

Table 1. Summary of the samples observed and breakdown of the classifications after follow-up spectroscopy

Candidate Sample		N	BCG & BCG?	Other ELGs & SA	QSO	Galaxies without emission	Stars	Not Classified
First priority	new	141	64	43	5	3	18	8
	already known	28	21	7	–	–	–	–
	total	169	85	50	5	3	18	8
Random sample	new	16	3	1	–	1	2	9
	already known	1	–	1	–	–	–	–
APM selected sample	new	22	10	10	–	–	1	1
	already known	1	–	1	–	–	–	–
Second priority	total	40	13	13	–	1	3	10
Objects presented in this paper		209	98	63	5	4	21	18

Table 2. Journal of observations

Date	Telescope	Instrument	Grating, grism	Wavelength range [Å]	Dispersion [Å/pixel]	Observed number
(1)	(2)	(3)	(4)	(5)	(6)	(7)
09.06-18.06.1999	2.2 m CAHA	CAFOS	G-200	3700–9500	4.5	117
—	—	—	B-200	3500–7400	4.7	18*
17.06-20.06.1999	4 m KPNO	R.C.Sp.	KPC-10A	3700–8300	2.8	46
08.12-11.12.1999	2.2 m CAHA	CAFOS	G-200	3700–9500	9.0	46

* – objects reobserved

The article is organized as follows. In section 2 we give the details of the spectroscopic observations and of the data reduction. In section 3 the results of the observations are presented in several tables. In section 4 we briefly discuss the new data and summarize the current state of the Hamburg/SAO survey. Throughout this paper a Hubble constant $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is used.

2. Spectral observations and data reduction

2.1. Observations

The results presented here were obtained in a snap-shot observing mode during two runs with the Calar Alto 2.2 m and one run with the KPNO 4 m telescopes (see Table 2).

2.2. Observations with the KPNO 4 m telescope

The observations were carried out with the Ritchey-Chretien Spectrograph attached to a Tektronix 2K×2K CCD detector. We used a $2'' \times 205''$ slit with a KPC-10A grating (316 grooves mm^{-1}) in its first order, and a GG 375 order separation filter cutting off second-order contamination for wavelengths blueward of 7400 Å. This instrumental setup allows a spatial scale along the slit of $0''.69 \text{ pixel}^{-1}$, a scale perpendicular to the slit of $2.77 \text{ Å pixel}^{-1}$, a spectral range of 3700 – 8300 Å and a spectral resolution of $\sim 7 \text{ Å}$ (FWHM). Short

exposures (3 – 5 minutes) were used in order to detect strong emission lines to allow measurement of redshifts and a crude classification. No orientation of the slit along the parallactic angle was done because of the snap-shot observing mode. Reference spectra of an Ar-Ne-He lamp were recorded to provide wavelength calibration. Spectrophotometric standard stars from Oke (1990) and Bohlin (1996) were observed at the beginning and at the end of each night for flux calibration. The dome flats, bias, dark and twilight sky frames were accumulated each night. The weather conditions were photometric, with seeing variations between $2''.5$ and $3''$ (FWHM).

2.3. Calar Alto 2.2 m telescope observations

Follow-up spectroscopy with the CAHA 2.2 m telescope was carried out during two runs (June and December, 1999, see Table 2), using the Calar Alto Faint Object Spectrograph (CAFOS) and Cassegrain focal reducer. During these runs a long slit of $300'' \times (2'' - 3'')$ and a G-200 grism (187 Å mm^{-1} , first order) were used. The B-200 grism (185 Å mm^{-1} , first order) was also used to reobserve 18 objects in order to improve the $[\text{OII}] \lambda 3727 \text{ Å}$ value. There were no order separation filters applied. The spatial scale along the slit was $0''.53 \text{ pixel}^{-1}$. A SITE 15 2K×2K CCD was operated in a 2×1 binned mode during the December run (binning only along the dispersion direc-

tion), while in the June run there was no binning applied. The wavelength coverages were $\lambda 3700 - \lambda 9500 \text{ \AA}$ with maximum sensitivity at $\sim 6000 \text{ \AA}$ for the G-200 grism and $\lambda 3500 - \lambda 7400 \text{ \AA}$ with maximum sensitivity at $\sim 4000 \text{ \AA}$ for the B-200 grism. The spectral resolution was $\sim 10 \text{ \AA}$ (FWHM) in the June run and due to CCD binning $\sim 20 \text{ \AA}$ (FWHM) in the December run. The slit orientation was not aligned with the parallactic angle because of the snapshot observing mode. The exposure times varied within 2–15 minutes depending on the object brightness. The observations were complemented by standard star flux measurements (Oke 1990, Bohlin 1996), reference spectra (Hg-Cd lamp) for wavelength calibration, dome flat, bias and dark frames. Most of the time the weather conditions were photometric with a seeing $\approx 1.5''$ (FWHM). Only during two nights the weather conditions were variable with a seeing of $3'' - 4''$ (FWHM). The measurements of these nights are marked by “*” in Table 4.

2.4. Data reduction

Reduction of the CAHA and KPNO spectral data was performed at the SAO using the standard reduction systems MIDAS¹ and IRAF².

The MIDAS command FILTER/COSMIC was found to be a quite successful way to remove automatically all cosmic ray hits from the images. After that we applied the IRAF package CCDRED for bad pixel removal, trimming, bias-dark subtraction, slit profile and flat-field corrections.

To do accurate wavelength calibration, correction for distortion and tilt for each frame, sky subtraction and correction for atmospheric extinction, the IRAF package LONGSLIT was used with invoking the IDENTIFY, REIDENTIFY, FITCOORD, TRANSFORM, BACKGROUND and EXTINCTION tasks.

To obtain an instrumental response function from observed spectrophotometric flux standards, the APSUM procedure from the APEXTRACT package was used first to extract apertures of standard stars. Then the sensitivity curve determined by the STANDARD and SENSFUNC procedures was applied by the CALIBRATE task to perform flux calibration for all object images. Finally the APSUM task was used to extract one-dimensional spectra from the flux calibrated images. In case that more than one exposure was obtained with the same setup for an object, the extracted spectra were co-added and a mean vector was calculated. In case of several observations with different setups (telescopes or grisms) for the same object, the data were reduced and measured independently and the more reliable values were taken.

¹ MIDAS is an acronym for the European Southern Observatory package – Munich Image Data Analysis System.

² IRAF is distributed by National Optical Astronomical Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation

To speed-up and facilitate the line measurements we employed dedicated command files created at the SAO using the FIT context and MIDAS command language. The procedures for the measurement of line parameters and redshifts applied were described in detail in Papers III and IV.

3. Results of follow-up spectroscopy

In Table 1 we present the results of the observations. The 209 candidates were selected from our first and second priority samples introduced in Paper IV.

Of 169 first priority candidates, 141 objects appeared in our list as new ones. 28 objects were listed in the NED as galaxies or objects from various catalogs with known redshifts and some of them already had information on emission lines in earlier publications. All objects were included in our observing program in order to improve spectral information. Comparison of our velocities with those of galaxies with already known redshift shows acceptable consistency within the uncertainties given.

Another 40 candidates observed were taken from the list of second priority candidates. As described in Paper IV two samples were created from this list. A “random selected sample” containing randomly selected objects from the list and the “APM selected sample” which uses additional information for selection. The “random selected sample” was created to access the fraction of BCGs in the second priority list and contains 43 objects. For 26 of them the spectral data were presented in Paper IV, while for the remaining 17 candidates observed with the CAHA 2.2 m telescope, the spectral information is presented here. The results of the analysis for this sample were presented in Paper IV. We found that the second priority list contains at most 10% BCG/HII-galaxies.

The second, “APM selected” sample comprises second priority candidates which are classified as non-stellar on Palomar Sky Survey plates (PSS) in the APM database, and have blue color according to the APM color system ($(B - R) < 1.0$). Here we give spectral data for 23 objects from this sample. Except one all are ELGs confirming the efficiency of this selection criterium to pick up the BCG/HII-galaxies from the second priority list (cf. Paper IV).

3.1. Emission-line galaxies

The observed emission line galaxies are listed in Table 3 containing the following information:

column 1: The object’s IAU-type name with the prefix HS.
column 2: Right ascension for equinox B1950.

column 3: Declination for equinox B1950. The coordinates were measured on direct plates of the HQS and are accurate to $\sim 2''$ (Hagen et al. 1995).

column 4: Heliocentric velocity and its r.m.s. uncertainty in km s^{-1} .

column 5: Apparent B-magnitude obtained by calibration of the digitized photoplates with photometric standard

stars (Engels et al. 1994), having an r.m.s. accuracy of $\sim 0^m.5$ for objects fainter than $m_B = 16^m.0$ (Popescu et al. 1996). Since the algorithm to calibrate the objective prism spectra is optimized for point sources the brightnesses of extended galaxies are underestimated. The resulting systematic uncertainties are expected to be as large as 2 mag (Popescu et al. 1996). For about 20% of our objects, B-magnitudes are unavailable at the moment. We present for them blue magnitudes obtained from the APM database. They are marked by a “+” before the value in the corresponding column. According to our estimate they are systematically brighter by $0^m.92$ than the B-magnitudes obtained by calibration of the digitized photoplates (r.m.s. $1^m.02$). Objects referred to Popescu & Hopp (2000) have precise B-magnitude in Vennik, Hopp & Popescu (2000). We do not list them here for the sake of homogeneity. The B-magnitude for HS 1213+3636A was determined by eye estimate as $17^m.5$ and marked by a “:” as less confident.

column 6: Absolute B-magnitude, calculated from the apparent B-magnitude and the heliocentric velocity. The only exception is made for HS 1213+3636B, a super-association in the nearby galaxy NGC 4214 for which the distance is known from stellar photometry (see comments below). No correction for galactic extinction is made because all objects are located at high galactic latitudes and the corrections are significantly smaller than the uncertainties of the magnitudes.

column 7: Preliminary spectral classification type according to the spectral data presented in this article. BCG means a galaxy possessing a characteristic HII-region spectrum with low enough luminosity ($M_B \geq -20^m$). SBN and DANS are galaxies of lower excitation with a corresponding position in line ratio diagrams, as discussed in Paper I. SBN are the brighter fraction of this type. Here we follow the notation of Salzer et al. (1989). Three objects were recognized as Seyfert galaxies. Two of them (HS 1317+4521B and HS 1616+3627) are Sy1 galaxies due to the presence of broad Balmer lines and of broad [FeII] emission. Our spectrum of HS 1616+3627 has insufficient quality to show this, but independent spectroscopy data of Grupe et al. (1999) clearly classify this object as a narrow-line Sy1 galaxy. The third one (HS 1220+3845) is a narrow-line ELG, which is classified as a Sy2 galaxy in diagnostic diagrams. Typical spectra of low-ionisation nuclear emission-line regions (LINERs) are identified for 3 galaxies. SA stands for two probable super-associations in two dwarf spiral galaxies. Six ELGs are difficult to classify. They are coded as NON.

column 8: One or more alternative names, according to the information from NED. References are given to the other sources of the redshift-spectral information indicating that a galaxy is an ELG.

The spectra of all emission-line galaxies are shown in Appendix A, which is available only in the electronic version of the journal.

The results of line flux measurements are given in Table 4. It contains the following information:

column 1: The object’s IAU-type name with the prefix

HS. By asterisk we note the objects observed during non-photometric conditions.

column 2: Observed flux (in 10^{-16} erg s $^{-1}$ cm $^{-2}$) of the H β line. For several objects without H β emission line the fluxes are given for H α and marked by a “+”.

columns 3,4,5: The observed flux ratios [OII]/H β , [OIII]/H β and H α /H β .

columns 6,7: The observed flux ratios [NII] λ 6583 Å/H α , and ([SII] λ 6716 Å + λ 6731 Å)/H α .

columns 8,9,10: Equivalent widths of the lines [OII] λ 3727 Å, H β and [OIII] λ 5007 Å. For several objects without detected H β emission line the equivalent widths are given for H α and marked by a “+”.

Below we give comments on some specific cases:

HS 0948+3723 and *HS 1430+4040:* H α -emission in these galaxies is affected by cosmic ray hits. The intensity ratio of H γ /H β is close to the recombination one with $c_{H\beta} = 0$. So we accepted for them the recombination H α /H β flux ratio, and corrected respectively the ratios of [NII]/H α and [SII]/H α in Table 4.

HS 1213+3636A: this is a compact slightly elongated ($\sim 9''$) object at $\sim 60''$ to NW from the center of the starbursting barred Magellanic type galaxy (SBm) NGC 4214 (with $M_B = -17^m.0$). Its radial velocity of 522 ± 35 km s $^{-1}$ is higher by ~ 220 km s $^{-1}$ than the systemic velocity of NGC 4214 and higher by ~ 260 km s $^{-1}$ than the velocity of H I gas in this place (McIntyre 1997). This implies that this compact emission region is a kinematically detached object, either a background dwarf, or more probably, a tiny “satellite” galaxy, passing by its much more massive neighbouring galaxy. For the former case its distance from the Hubble flow is 7 Mpc, $M_B = -11^m.7$; and the size ~ 0.3 kpc. For the latter case, if HS 1213+3636A is located at the same distance as NGC 4214 (2.64 Mpc, as found through stellar photometry with HST by Drozdovsky et al. 2001), its $M_B = -9^m.6$; and the size ~ 0.1 kpc. The nature of this tiny galaxy can be checked in principle by the use of well known luminosity-abundance relationship for gas-rich galaxies. If it is situated far from NGC 4214 one should expect that this galaxy should have metallicity $Z \leq 1/30 Z_\odot$. On the other hand, if HS 1213+3636A is connected to NGC 4214, one can speculate that this companion formed of gas pulled out of the parent SBm galaxy as a result of its strong interaction with other galaxy. In this case the metallicity of this companion should be close to that of NGC 4214, that is known to be $\sim 1/4 - 1/5 Z_\odot$ (e.g., Kobulnitsky & Skillman 1996). Thus, metallicity determination of this dwarf with strong starburst will probably resolve the dilemma.

HS 1213+3636B: this well elongated object ($\sim 12''$) is projected onto the middle part ($\sim 46''$, or ~ 0.6 kpc to NWW from the center) of the same galaxy NGC 4214. It is at $\sim 25''$ to the South from HS 1213+3636A. Our velocity differs by only ~ 60 km s $^{-1}$ from the NGC 4214 systemic velocity and is well consistent with the H I velocity of 260 km s $^{-1}$ at this place from McIntyre (1997). From our data it follows that HS 1213+3636B is a

super-association (SA) in a spiral arm of NGC 4214. Its $M_B = -9^m8$: and the size ~ 0.15 kpc. The spectra of both A and B objects were acquired in one slit, so their large velocity difference is out of doubt.

HS 1311+3628: this elongated object ($\sim 11''$) is projected onto the southern edge ($\sim 50''$ to the South from the center) of Im galaxy UGC 8303 (Holmberg VIII). A large difference of its radial velocity (1094 ± 12 km s $^{-1}$) with that of the systemic velocity of the dwarf irregular UGC 8303 (944 ± 5 km s $^{-1}$ from Huchra et al. 1995) and the local HI velocity in this region (~ 945 km s $^{-1}$) from the HI map by Thean et al. (1997) (where this galaxy by error is called UGC 8314), implies that this object probably does not belongs to UGC 8303 and is a nearby separate star-forming dwarf. We observed this object twice on different nights and with different setups. Both spectra have the same redshifts within the observational uncertainties. HS 1311+3628 is therefore not an HII-region in UGC 8303 as suggested by Popescu et al. (1996).

HS 1311+3924: this is the irregular (probably merging) galaxy UGC 8315 and member of a group, with a radial velocity of 1215 km s $^{-1}$ according to Garcia et al. (1993). We got the spectrum of only one of two bright knots in the galaxy elongated body, namely the NE one. While its velocity 1401 ± 43 differs by 186 km s $^{-1}$ from that given by Garcia et al. (1993) the combined uncertainty of latter value is not well constrained. A long-slit spectrum along the major axis will be helpful for understanding the nature of this object.

HS 1423+3945: the radial velocity of this object is very close to that of UGC 9242 (our 1480 ± 35 versus 1440 ± 11 km s $^{-1}$ in the RC3 catalog) and therefore the object is very probably an SA in this edge-on dwarf spiral.

HS 1242+4058A: The nearby BCG HS 1242+4058B (cf. Paper II) is separated by $\sim 4'$ from HS 1242+4058A and the radial velocities differ by ≈ 300 km/s. They are probably not associated with each other.

HS 1317+4521B: The nearby BCG-candidate ($\sim 10''$ to SW) HS 1317+4521A is not observed yet.

HS 1646+4003B: its faint companion ($\sim 30''$ to SSW), HS 1646+4003A was earlier identified as NON (cf. Paper II).

HS 0935+4135, *HS 0958+3654*, *HS 1025+3707B*, *HS 1027+4728*, *HS 1206+4012*, *HS 1236+4532*, *HS 1433+4245*, *HS 1514+4602*: In all these galaxies the only emission line suitable for redshift measurement is H α . A conservative estimate of their radial velocity uncertainty is 300 km s $^{-1}$.

For some of the galaxies either the snap-shot spectra had strong enough emission lines and a well detected [OIII] $\lambda 4363$ Å or they were reobserved later with better S/N ratio. For these galaxies we determined metallicities and the following galaxies appeared to have $O/H \leq 1/10$ (O/H) $_{\odot}$: HS 1313+4521, HS 1330+3651, HS 1334+3957, HS 1408+4227, HS 1417+4433, HS 1545+4055, HS 1650+3706, HS 1655+3845.

For two BCGs (HS 0940+4052 and HS 1311+3628) the blue WR-bump is more or less evident from the snap-shot spectra, and for 8 more BCGs this feature is probably present.

Several “BCG?” were listed in Table 3 with absolute magnitudes brighter than -20^m0 . In fact, these M_B are uncertain, since they come from the estimates from the Hamburg Quasar Survey plates. Independent CCD measurements or estimates from alternative sources show that usually they are fainter by $0^m5 - 1^m0$, and should fulfill therefore the criterion $M_B \geq -20^m0$ (Kniazhev et al. 2001b).

3.2. Quasars

The main criteria applied to search for BCGs are a blue continuum near $\lambda 4000$ Å and a strong emission line — expected is [OIII] $\lambda 5007$ Å — in the wavelength region between 5000 Å and the sensitivity break of the Kodak IIIa-J photoemulsion near 5400 Å (see Paper I). For this reason faint QSOs with Ly α $\lambda 1216$ Å redshifted to $z \sim 3$ or with MgII $\lambda 2798$ Å redshifted to $z \sim 0.8$ could be selected as BCG candidates. In Papers I–IV we already reported on the discovery of a number of such faint QSOs. They have been missed by the proper Hamburg Quasar Survey since the latter is restricted to bright QSOs ($B \leq 17 - 17.5$). Here we report on the discovery of five high-redshift faint ($B > 17.5$) QSOs. For all but one of them, we identified Ly α $\lambda 1216$ Å redshifted to $z \sim 3$ as the line responsible for its selection. One object (HS 1301+4233) shows a broad emission line tentatively identified with MgII $\lambda 2798$ Å. However this identification remains somewhat uncertain due to low signal-to-noise of the spectrum. Two of our new QSOs: HS 1224+4410 and HS 1541+4452, have recently appeared in the NED as radio sources with no redshift determination (reported as [HVG99] R17 in Hunter et al. (1999) and B3 1541+448A in Douglas et al. (1996)). The data for all five quasars are presented in Table 5. Finding charts and plots of their spectra can be found on the www-site of the Hamburg Quasar Survey (<http://www.hs.uni-hamburg.de/hqs.html>).

3.3. Non-emission-line objects

In total, for 43 candidates no (trustworthy) emission lines were detected. We divided them into three categories.

3.3.1. Absorption-line galaxies

For four non-ELG galaxies the signal-to-noise ratio of our spectra was sufficient to detect absorption lines, allowing the determination of redshifts. The data are presented in Table 6.

3.3.2. Stellar objects

To separate the stars among the objects missing detectable emission lines we cross-correlated a list of the most common stellar features with the observed spectra. In total, 21 objects with definite stellar spectra and redshifts close to zero were identified. All of them were crudely classified in categories from definite A-stars to G-stars, with most of them intermediate between A and F. The data for these stars are presented in Table 7.

3.3.3. Non-classified objects

There was no possibility of classifying 18 objects without emission lines. Their spectra have too low signal-to-noise ratio to detect trustworthy absorption features, or the EWs of their emission lines are too small.

4. Discussion

4.1. The fifth list

As result we have 209 observed candidates preselected on HQS objective prism plates, of which 169 were first priority candidates and 40 were second priority ones. 166 objects (79 % from 209 objects) are found to be either ELGs (161), or quasars (5).

Of 161 ELGs 98 galaxies (61 %) were classified based on the character of their spectra and their luminosity as HII/BCGs or probable BCGs.

Two of them (HS 1213+3636A and HS 1311+3628) are low-mass neighbours/satellites of the dwarf spiral NGC 4214 and of the Im galaxy UGC 8303. As the discussion of the local environment of BCGs is out of the scope of this paper, we refer to a recent analysis of this issue by Pustilnik et al. (2001).

HS 1213+3636B and HS 1423+3945 are SAs in the dwarf spirals NGC 4214 and UGC 9242.

Six ELGs are difficult to classify at all due to their poor signal-to-noise spectra. Six more ELGs were classified as Active Galactic Nuclei (AGNs): 3 as Seyfert galaxies and 3 as LINERs. The remaining 49 ELGs are objects with low excitation: either starburst nuclei galaxies (SBN and probable SBN) or their lower mass analogs – dwarf amorphous nuclear starburst galaxies (DANS or probable DANS).

By keeping a high fraction of BCGs (~ 62 % among the first priority candidates) we continue to have a high efficiency of discovery new BCGs, which is the main goal of the HSS. Since the completeness of the BCG sample under construction is an important parameter for many follow-up statistical studies, we observed a randomly selected sample of candidates from our list of second priority candidates. As discussed already in Paper IV at most 10 % of them turned out to be BCGs. Using additional information from the APM is an efficient means to pick up these BCGs among the second priority candidates: among 23 objects observed 10 BCGs (43 %) were discovered (Table 1).

4.2. Summary of the present status of the survey

Summarizing the results of the Hamburg/SAO survey presented in Papers I through V, we discovered altogether from the 1-st priority candidates 433 new emission-line objects (25 of them are QSOs), and for 85 known ELGs we got quantitative data for their emission lines. At the moment the total number of confident or probable blue compact/low-mass HII-galaxies reaches 360. Relative to all observed 493 ELGs the fraction of BCGs (360/493 or 73 %) demonstrates the high efficiency of the survey to find this type of galaxies. 21 more new BCGs and 20 other type ELGs are found among the second priority candidates. To estimate the total number of BCGs in the HSS zone we should count new BCGs expected from the remaining candidates and those selected in the HSS, but not observed by us since they already were known from other surveys. Thus we expect the total number of BCGs in this sky region to be ~ 500 . This will be the largest homogeneous BCG sample in both hemispheres.

5. Conclusions

We made follow-up spectroscopy of the fifth list of candidates from the Hamburg/SAO Survey for ELGs. Summarizing the results of the spectroscopy, the analysis of spectral information and the discussion above we draw the following conclusions:

- The intended methods to detect ELG candidates on the plates of the Hamburg Quasar Survey give a reasonably high detection rate of emission-line objects. In total, out of both priority categories, we observed 209 objects among which we found 166 emission-line objects corresponding to a detection rate of ~ 79 %.
- Besides the ELGs we found also 5 new quasars, mostly with $\text{Ly}\alpha$ in the wavelength region 4950 – 5100 Å ($z \sim 3$), near the red boundary of the IIIa-J photo-plates.
- The high fraction of BCG/HII galaxies among all observed ELGs (about 61 % in this paper) is in line with our main goal – to pick up efficiently a statistically well selected and deep BCG sample in the sky region under analysis.

Acknowledgements. This work was supported by the grant of the Deutsche Forschungsgemeinschaft No. 436 RUS 17/77/94. U.A.V. is very grateful to the staff of the Hamburg Observatory for their hospitality and kind assistance. Support by the INTAS grant No. 96-0500 was crucial to proceed with the Hamburg/SAO survey declination band centered on $+37.5^\circ$. SAO authors acknowledge also partial support from the INTAS grant No. 97-0033. We note that the use of APM facility was extremely valuable for selection methodology of additional candidates to BCGs from the 2-nd priority list. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. We have also used the Digitized Sky Survey, produced at the Space Telescope Science Institute under government grant NAG W-2166.

References

- Bohlin, R. C. 1996, *AJ*, 111, 1743
- Douglas, J. N., Bash, F. N., Bozayan, F. A., Torrence, G. W., & Wolfe, C. 1996, *AJ*, 111, 1945
- Drozdovsky, I., et al. 2001, in Bad Honnef conference “Dwarf galaxies and their environment”, Jan. 23–27, 2001
- Engels, D., Cordis, L., & Köhler, T. 1994, *Proc. IAU Symp.* 161, ed. H. T. MacGillivray (Kluwer: Dordrecht), 317
- Garcia, A. M., Bottinelli, L., Garnier, R., Gouguenheim, L., & Paturel, G. 1993, *A&AS*, 97, 801
- Grupe, D., Beuermann, K., Mannheim, K., & Thomas H.-C. 1999, *A&A*, 350, 895
- Hagen, H.-J., Groote, D., Engels, D., & Reimers, D. 1995, *A&AS*, 111, 195
- Hopp, U., Engels, D., Green, R., et al. 2000, *A&AS*, 142, 417 (**Paper III**)
- Huchra, J. P., Geller, M. J., & Corwin, H. G. Jr. 1995, *ApJS*, 99, 391
- Hunter, D., van Woerden, H., & Gallagher, J. 1999, *AJ*, 118, 2184
- Kniazev, A. Y., Engels, D., Pustilnik, S. A. et al. 2001a, *A&A*, 366, 771 (**Paper IV**)
- Kniazev, A. Y., Pustilnik, S. A., Ugryumov A.V. et al. 2001b, in preparation
- Kobulnicky, H.A., Skillman, E.D. 1996, *ApJ*, 471, 211
- Markarian, B. E., Lipovetsky, V. A., & Stepanian, J. A. 1983, *Afz*, 19, 29
- McIntyre, V. J. 1997, *Publ. Astron. Soc. Australia*, 14, 122
- Oke, J. B. 1990, *AJ*, 99, 1621
- Pesch, P., Stephenson, C. B., & MacConnell, D. J. 1995, *ApJS*, 98, 41
- Popescu, C. C., & Hopp, U. 2000, *A&AS*, 142, 247
- Popescu, C. C., Hopp, U., Hagen, H.-J., & Elsässer, H. 1996, *A&AS*, 116, 43
- Pustilnik, S. A., Engels, D., Ugryumov, A. V., et al. 1999, *A&AS*, 135, 299 (**Paper II**)
- Pustilnik, S. A., Kniazev, A. Y., Lipovetsky, V. A., & Ugryumov, A. V. 2001, *A&A*, in press = astro-ph/0104334
- Salzer, J. J., MacAlpine, G. M., & Boroson, T. A. 1989, *ApJS*, 70, 479
- Salzer, J. J., Moody, J. W., Rosenberg, J. L., Gregory, S. A., & Newberry, M. V. 1995, *AJ*, 109, 2376
- Stepanian, J. A. 1994, *Proc. IAU Symp.* 161, ed. H. T. MacGillivray (Kluwer: Dordrecht), 731
- Thean, A. H. C., Mundell, C. G., Pedlar, A., & Nicholson, R. A. 1997, *MNRAS*, 290, 15
- Ugryumov, A. V., Pustilnik, S. A., Lipovetsky, V. A., Izotov, Yu. I., & Richter, G. M. 1998, *A&AS*, 131, 295
- Ugryumov, A. V., Engels, D., Lipovetsky, V. A., et al. 1999, *A&AS*, 135, 511 (**Paper I**)
- Vennik, J., Hopp, U., & Popescu, C. C. 2000, *A&AS*, 142, 399

Table 3. Coordinates, Velocities and Magnitudes of Emission-Line Galaxies

#	Name	α (1950)	δ (1950)	$V_{hel}^a \pm \sigma_V$	m_B	M_B^b	Type	Other names from NED and number of reference
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	
1	HS 0809+4334	08 09 01.5	+43 34 17	14146 $\pm$ 57	18.4	-18.0	BCG	
2	HS 0818+4234	08 18 40.5	+42 34 23	17424 $\pm$ 68	19.7	-17.1	NON	
3	HS 0822+4049	08 22 25.1	+40 49 57	12087 $\pm$ 100	17.3	-18.7	BCG	
4	HS 0822+4238	08 22 50.9	+42 38 24	11637 $\pm$ 60	18.2	-17.8	BCG	
5	HS 0824+4139	08 24 11.6	+41 39 00	10784 $\pm$ 53	18.4	-17.4	BCG?	IRAS F08241+4139
6	HS 0828+4305	08 28 04.6	+43 05 24	10700 $\pm$ 86	16.7	-19.0	SBN	
7	HS 0828+4346	08 28 32.0	+43 46 24	7649 $\pm$ 62	18.4	-16.6	BCG	
8	HS 0831+4208	08 31 14.5	+42 08 17	44859 $\pm$ 56	+17.3	-21.5	SBN	MBWU023
9	HS 0849+3802 [‡]	08 49 09.9	+38 02 27	13368 $\pm$ 71	16.9	-19.4	SBN	IRAS F08491+3802
10	HS 0849+3639 [‡]	08 49 34.0	+36 39 14	7503 $\pm$ 47	20.0	-15.0	BCG	
11	HS 0850+3722 [‡]	08 50 17.5	+37 22 40	14925 $\pm$ 68	+18.7	-17.8	NON	
12	HS 0855+4118	08 55 02.3	+41 18 44	8113 $\pm$ 80	17.0	-18.2	BCG?	
13	HS 0855+3729 [‡]	08 55 27.6	+37 29 06	14921 $\pm$ 59	17.5	-19.0	SBN	
14	HS 0900+4318	09 00 26.4	+43 18 52	2421 $\pm$ 95	18.5	-14.0	BCG	MBWU369
15	HS 0901+3517 [‡]	09 01 17.6	+35 17 29	12397 $\pm$ 95	18.2	-17.9	BCG?	
16	HS 0904+3752 [‡]	09 04 29.2	+37 52 27	6803 $\pm$ 86	18.4	-16.4	DANS?	FGC0841
17	HS 0907+4201	09 07 43.6	+42 01 48	12004 $\pm$ 104	17.1	-18.9	SBN	
18	HS 0912+4427	09 12 08.2	+44 27 26	11637 $\pm$ 101	17.4	-18.6	DANS?	
19	HS 0912+4259	09 12 23.6	+42 59 50	14268 $\pm$ 53	16.4	-20.0	SBN	IRAS F09123+4259
20	HS 0912+4433	09 12 28.7	+44 33 55	11345 $\pm$ 92	17.3	-18.6	BCG	
21	HS 0912+3900 [‡]	09 12 38.2	+39 00 21	20628 $\pm$ 92	17.2	-20.0	SBN	NPM1G+39.0196
22	HS 0923+4021	09 23 13.8	+40 21 32	5190 $\pm$ 110	17.2	-17.0	BCG	
23	HS 0927+4159	09 27 00.6	+41 59 35	13533 $\pm$ 95	17.4	-18.9	BCG	
24	HS 0934+4150	09 34 21.3	+41 50 29	12011 $\pm$ 74	17.0	-19.0	BCG	KUG0934+418
25	HS 0934+4256	09 34 53.9	+42 56 20	3600 $\pm$ 44	17.2	-16.2	DANS	KUG0934+429
26	HS 0935+4135	09 35 30.7	+41 35 07	79288 $\pm$ 300	18.0	-22.1	LINER	
27	HS 0938+4215	09 38 49.6	+42 15 40	4680 $\pm$ 98	17.6	-16.4	BCG	
28	HS 0939+4124	09 39 37.2	+41 24 03	13646 $\pm$ 62	18.5	-17.8	BCG	
29	HS 0940+4052	09 40 08.1	+40 52 28	18255 $\pm$ 44	16.1	-20.8	BCG?	
30	HS 0941+4024A	09 41 54.7	+40 24 52	21709 $\pm$ 20	+17.2	-20.1	SBN	FIRSTJ094501.7+40110
31	HS 0941+4024B	09 41 56.2	+40 24 53	21724 $\pm$ 14	16.6	-20.7	BCG?	
32	HS 0947+3526 [‡]	09 47 59.6	+35 26 39	4678 $\pm$ 44	16.4	-17.6	BCG	KUG0947+354
33	HS 0948+3723	09 48 19.0	+37 23 05	5140 $\pm$ 37	+17.8	-16.4	BCG	6
34	HS 0951+3712 [‡]	09 51 40.6	+37 12 07	12275 $\pm$ 65	16.9	-19.2	SBN	FIRSTJ095440.9+36575
35	HS 0952+4024 [‡]	09 52 30.8	+40 24 28	8648 $\pm$ 50	17.3	-18.0	DANS	
36	HS 0953+3628	09 53 16.5	+36 28 41	14435 $\pm$ 101	17.3	-19.1	BCG	CG 308, 5
37	HS 0954+3615 [‡]	09 54 35.4	+36 15 47	5193 $\pm$ 68	18.4	-15.8	BCG	
38	HS 0956+3702 [‡]	09 56 08.4	+37 02 05	15679 $\pm$ 74	16.7	-19.9	SBN	
39	HS 0958+3654 [‡]	09 58 27.9	+36 54 43	12265 $\pm$ 300	17.4	-18.7	SBN?	CGCG182-058, 8
40	HS 0958+3637	09 58 29.2	+36 37 30	13134 $\pm$ 75	17.6	-18.6	DANS?	

Table 3. (Continued)

#	Name	α (1950)	δ (1950)	$V_{hel}^a \pm \sigma_V$	m_B	M_B^b	Type	Other names from NED and number of reference
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
41	HS 0958+3718	09 58 15.5	+37 18 38	1606 ± 77	15.7	-16.0	BCG	CG 315,KUG0958+373, 5
42	HS 1004+3730 [‡]	10 04 22.3	+37 30 22	10133 ± 94	16.5	-19.2	SBN	KUG1004+375
43	HS 1008+3620 [‡]	10 08 01.5	+36 20 51	11424 ± 53	17.2	-18.7	BCG	
44	HS 1011+3604	10 11 29.6	+36 04 55	10227 ± 65	17.5	-18.2	DANS?	
45	HS 1014+3535 [‡]	10 14 50.0	+35 35 50	26718 ± 72	17.6	-20.2	SBN	KUG1014+355
46	HS 1016+3733	10 16 02.7	+37 33 04	14451 ± 78	16.1	-20.3	LINER	KUG1016+375
47	HS 1021+4839	10 21 39.0	+48 39 53	7495 ± 69	16.7	-18.3	SBN?	
48	HS 1025+3707A	10 25 12.7	+37 07 08	7503 ± 102	17.9	-17.1	BCG	
49	HS 1025+3707B	10 25 13.7	+37 07 10	39650 ± 300	+16.9	-21.7	SBN	
50	HS 1025+3658 [‡]	10 25 32.3	+36 58 28	10988 ± 59	16.9	-18.9	BCG	
51	HS 1025+4057	10 25 51.0	+40 57 52	16482 ± 59	16.7	-20.0	SBN	
52	HS 1027+4728	10 27 45.9	+47 28 57	18225 ± 100	17.6	-19.3	SBN	
53	HS 1028+3716	10 28 24.8	+37 16 15	11137 ± 109	16.9	-19.0	DANS	KUG1028+372
54	HS 1032+4600	10 32 21.2	+46 00 11	9299 ± 61	17.1	-18.4	BCG?	KUG1032+460
55	HS 1043+3852 [‡]	10 43 09.0	+38 52 15	10071 ± 60	+18.9	-16.7	BCG	
56	HS 1122+3800 [‡]	11 22 53.7	+38 00 33	11438 ± 98	+18.7	-17.2	BCG	
57	HS 1148+3732	11 48 05.4	+37 32 22	10835 ± 51	17.5	-18.3	DANS	KUV11481+3732, 14
58	HS 1159+3940	11 59 52.5	+39 40 59	6299 ± 29	+17.4	-17.2	DANS	6
59	HS 1200+3824	12 00 41.7	+38 24 01	23983 ± 146	+18.0	-19.4	SBN?	MAPS-NGPO 266 006267
60	HS 1201+3535	12 01 17.6	+35 35 08	10216 ± 11	19.0	-16.7	BCG	MAPS-NGPO 267 026327
61	HS 1202+3732	12 02 36.8	+37 32 55	6787 ± 7	+18.7	-16.1	BCG	MAPS-NGPO 267 0065953
62	HS 1204+3945	12 04 45.2	+39 45 47	11178 ± 195	18.0	-17.9	BCG?	
63	HS 1205+3940	12 05 29.2	+39 40 19	6362 ± 14	18.1	-16.5	BCG	CG 1517
64	HS 1206+4012	12 06 19.8	+40 12 24	26721 ± 300	18.1	-19.7	NON	
65	HS 1210+3831	12 10 46.7	+38 31 42	15800 ± 66	16.9	-19.7	BCG?	CG 1538
66	HS 1211+3807	12 11 20.1	+38 07 27	7433 ± 133	16.3	-18.7	BCG	MAPS-NGPO 267 004482
67	HS 1211+4012	12 11 51.5	+40 12 23	7091 ± 23	18.3	-16.6	BCG	
68	HS 1212+3640	12 12 09.5	+36 40 57	12704 ± 56	16.5	-19.6	BCG	KUG1212+366
69	HS 1213+3636A	12 13 03.8	+36 36 56	522 ± 35	17.5:	-11.7:	BCG	
70	HS 1213+3636B	12 13 04.1	+36 36 31	229 ± 38	17.3	-9.6	SA in SBm	in NGC 4214
71	HS 1215+3850	12 15 46.7	+38 50 07	15702 ± 65	16.1	-20.5	BCG?	CG 1553
72	HS 1216+3743	12 16 38.5	+37 43 24	5978 ± 124	17.9	-16.6	DANS	MAPS-NGPO 267 007023
73	HS 1218+3945	12 18 52.5	+39 45 44	11475 ± 57	17.7	-18.2	BCG	
74	HS 1220+3845	12 20 11.8	+38 45 38	57118 ± 57	+17.8	-21.6	Sy2	
75	HS 1223+3938	12 23 29.6	+39 38 31	10873 ± 69	16.8	-19.0	BCG	NPM1G+39.0289,CG 1566, 2,7
76	HS 1224+4027	12 24 42.2	+40 27 56	17724 ± 5	18.6	-18.3	BCG	
77	HS 1225+4129	12 25 45.5	+41 29 16	11411 ± 17	18.7	-17.2	BCG	
78	HS 1226+4113	12 26 55.8	+41 13 52	3845 ± 17	18.7	-14.9	BCG	
79	HS 1232+3834	12 32 25.0	+38 34 31	4049 ± 57	+16.8	-16.9	BCG	MAPS-NGPO 218 110114
80	HS 1232+3947	12 32 54.6	+39 47 39	6506 ± 143	17.2	-17.5	BCG	2,7
81	HS 1235+4232	12 35 13.7	+42 32 02	7449 ± 90	18.4	-16.6	DANS?	
82	HS 1236+3821	12 36 31.9	+38 21 52	2069 ± 103	15.4	-16.8	BCG	UGC 7816,CGCG1236.5+3822, NPM1G+38.0259, 2,7

Table 3. (Continued)

#	Name	α (1950)	δ (1950)	$V_{hel}^a \pm \sigma_V$	m_B	M_B^b	Type	Other names from NED and number of reference
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
83	HS 1236+4532	12 36 39.8	+45 32 54	16334 $\pm$ 300	17.6	-19.1	NON	MAPS-NGPO 172 027192
84	HS 1237+3900	12 37 04.0	+39 00 00	11198 $\pm$ 11	+18.6	-17.3	BCG	6
85	HS 1238+4229	12 38 24.1	+42 29 08	11370 $\pm$ 53	18.9	-17.0	BCG	
86	HS 1242+4058A	12 42 00.3	+40 58 12	5512 $\pm$ 197	16.6	-17.7	BCG	MAPS-NGPO 218 061180
87	HS 1244+3647	12 44 36.9	+36 47 45	14171 $\pm$ 89	16.4	-20.0	BCG	2,7
88	HS 1245+4413	12 45 52.8	+44 13 25	9293 $\pm$ 62	17.7	-17.8	DANS	
89	HS 1300+4225	13 00 36.5	+42 25 39	11374 $\pm$ 23	+17.3	-18.6	BCG?	
90	HS 1302+4058	13 02 13.6	+40 58 51	9200 $\pm$ 34	+16.8	-18.6	BCG	MAPS-NGPO 219 053488
91	HS 1302+4101	13 02 49.8	+41 01 22	10567 $\pm$ 20	+18.5	-17.2	BCG	
92	HS 1305+4049	13 05 49.2	+40 49 09	10456 $\pm$ 57	+17.8	-17.9	BCG	
93	HS 1306+4146	13 06 13.5	+41 46 23	17237 $\pm$ 5	+19.1	-17.7	BCG	
94	HS 1306+4151	13 06 46.9	+41 51 04	82269 $\pm$ 23	+18.0	-22.2	NON	
95	HS 1306+4232	13 06 57.7	+42 32 19	14701 $\pm$ 90	+16.9	-19.6	DANS	MAPS-NGPO 219 019849
96	HS 1307+4225	13 07 38.8	+42 25 04	2677 $\pm$ 20	+17.4	-15.3	BCG	MAPS-NGPO 219 022840
97	HS 1309+4446	13 09 20.6	+44 46 42	8944 $\pm$ 158	+16.4	-19.0	DANS	
98	HS 1310+4224	13 10 20.8	+42 24 08	12565 $\pm$ 117	+17.7	-18.4	DANS	
99	HS 1311+3628	13 11 00.0	+36 28 02	1094 $\pm$ 12	18.0	-12.9	BCG	Near UGC 8303, 2,7 NGP9 F269-0669419 is 14'' to E
100	HS 1311+4507	13 11 30.2	+45 07 33	18285 $\pm$ 11	+18.6	-18.4	BCG	
101	HS 1311+4204	13 11 50.0	+42 04 50	10495 $\pm$ 14	+18.1	-17.6	DANS	
102	HS 1311+3924	13 11 54.2	+39 24 47	1401 $\pm$ 54	15.6	-15.8	BCG	UGC 8315, 4
103	HS 1313+4440	13 13 04.4	+44 40 15	10982 $\pm$ 144	16.5	-19.3	SBN	UGC 8327,MRK 248, 9
104	HS 1313+4246	13 13 25.5	+42 46 04	12055 $\pm$ 11	+19.2	-16.9	BCG	MAPS-NGPO 219 017637
105	HS 1313+4041	13 13 34.2	+40 41 07	6373 $\pm$ 23	+17.6	-17.1	BCG	
106	HS 1313+4521	13 13 40.2	+45 21 59	3449 $\pm$ 18	+18.9	-14.4	BCG	
107	HS 1314+4145	13 14 00.0	+41 45 28	6445 $\pm$ 127	15.4	-19.3	SBN	MRK 1477, 10
108	HS 1315+4129	13 15 42.5	+41 29 14	35297 $\pm$ 116	+18.0	-20.3	NON	FIRSTJ131755.8+41132, 7C1315+4129
109	HS 1316+3839	13 16 01.7	+38 39 22	18065 $\pm$ 86	16.4	-20.5	BCG?	
110	HS 1317+4521B	13 17 50.9	+45 21 60	32888 $\pm$ 133	+16.3	-21.9	Sy1	
111	HS 1322+3708 [†]	13 22 17.0	+37 08 39	14127 $\pm$ 67	16.4	-20.0	BCG	KUG1322+371
112	HS 1323+3946	13 23 29.9	+39 46 13	7064 $\pm$ 162	16.0	-18.9	DANS	IRAS F13235+3945
113	HS 1330+3651	13 30 54.0	+36 51 55	5044 $\pm$ 117	16.7	-17.4	BCG	2,7
114	HS 1334+3957	13 34 11.4	+39 57 31	2513 $\pm$ 61	16.9	-15.7	BCG	2,7
115	HS 1336+3650	13 36 43.9	+36 50 57	5791 $\pm$ 94	17.8	-16.6	BCG	2,7
116	HS 1354+3635	13 54 29.9	+36 35 39	5056 $\pm$ 56	16.1	-18.0	BCG	CG 1201, 3,2,7
117	HS 1402+3650	14 02 39.2	+36 50 47	10456 $\pm$ 83	16.3	-19.4	BCG	2,7
118	HS 1408+4227	14 08 07.5	+42 27 26	20355 $\pm$ 17	20.6	-16.6	BCG	
119	HS 1408+4522	14 08 09.3	+45 22 23	20443 $\pm$ 113	17.8	-19.4	BCG	CG 367
120	HS 1410+3627	14 10 02.2	+36 27 15	9890 $\pm$ 90	16.9	-18.7	BCG	2,7
121	HS 1412+4203	14 12 12.3	+42 03 50	24842 $\pm$ 160	19.1	-18.5	SBN	
122	HS 1414+4034	14 14 43.1	+40 34 58	6097 $\pm$ 32	20.1	-14.5	BCG	
123	HS 1417+4433	14 17 27.9	+44 33 14	18797 $\pm$ 5	19.8	-17.2	BCG	

Table 3. (Continued)

#	Name	α (1950)	δ (1950)	$V_{hel}^a \pm \sigma_V$	m_B	M_B^b	Type	Other names from NED and number of reference
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
124	HS 1419+3639	14 19 16.2	+36 39 18	3579 ± 42	17.1	-16.3	BCG	UGC 9198, 2
125	HS 1420+3726 [†]	14 20 50.1	+37 26 58	10847 ± 45	17.5	-18.3	BCG	
126	HS 1423+3945 [†]	14 23 13.6	+39 45 27	1480 ± 35	18.5	-13.0	SA in Sp	in UGC 9242
127	HS 1423+4054	14 23 50.0	+40 54 48	8059 ± 93	18.5	-16.7	BCG	
128	HS 1427+4309	14 27 53.4	+43 09 40	2679 ± 151	17.8	-15.0	BCG	CG 449,PGC051797, [KOS87]142752+430936, 12
129	HS 1428+4056	14 28 15.9	+40 56 16	10276 ± 49	19.9	-15.8	DANS	
130	HS 1430+4040	14 30 17.9	+40 40 54	6999 ± 60	17.2	-17.7	DANS	CG 467,IRAS F14302+4041, PGC051943, 13
131	HS 1433+4411	14 33 27.2	+44 11 21	2945 ± 90	16.8	-16.2	BCG	CG 480,NPM1G+44.0284
132	HS 1433+4245	14 33 53.6	+42 45 23	21481 ± 300	14.0	-23.3	LINER	NPM1G+42.0387
133	HS 1437+4002	14 37 58.5	+40 02 29	11084 ± 29	+19.2	-16.7	BCG	6
134	HS 1441+4122	14 41 24.4	+41 22 44	11007 ± 122	17.3	-18.5	DANS	NPM1G+41.0365
135	HS 1443+4027	14 43 11.0	+40 27 39	8425 ± 30	17.1	-18.2	BCG	CG 527
136	HS 1447+4402	14 47 20.1	+44 02 50	24642 ± 56	19.2	-18.4	DANS	
137	HS 1448+4518	14 48 56.3	+45 18 01	10661 ± 149	17.8	-18.0	DANS?	
138	HS 1450+3844	14 50 20.8	+38 44 37	4179 ± 54	17.8	-15.9	BCG	CG 565,[SP82]25, 2
139	HS 1450+4234	14 50 55.3	+42 34 20	10393 ± 194	17.4	-18.3	DANS	CG 568,NPM1G+42.0398
140	HS 1458+4129	14 58 03.9	+41 29 17	8687 ± 116	18.4	-16.9	DANS	
141	HS 1514+4602	15 14 27.6	+46 02 48	22810 ± 300	16.8	-20.7	SBN	NPM1G+46.0322, 1
142	HS 1537+4411	15 37 19.2	+44 11 23	9312 ± 43	17.2	-18.3	BCG	
143	HS 1545+4055	15 45 26.2	+40 55 36	9521 ± 14	18.8	-16.7	BCG	[SP82]54
144	HS 1545+4459	15 45 34.0	+44 59 19	10480 ± 114	18.4	-17.3	BCG?	
145	HS 1546+4319	15 46 31.9	+43 19 50	12400 ± 45	16.5	-19.6	SBN	IRAS F15465+4320
146	HS 1603+3759 [‡]	16 03 13.2	+37 59 08	12071 ± 82	18.6	-17.4	BCG	
147	HS 1613+3902	16 13 51.4	+39 02 28	10046 ± 77	18.7	-16.9	BCG	FIRSTJ161538.5+38544
148	HS 1616+3627 [†]	16 16 19.9	+36 27 11	10210 ± 38	16.5	-19.2	Sy1	RXJ1618.1+3619, 11
149	HS 1625+3847	16 25 49.1	+38 47 28	9355 ± 131	18.4	-17.1	BCG?	
150	HS 1633+4703	16 33 12.8	+47 03 44	2681 ± 62	16.7	-16.1	BCG	2,7
151	HS 1645+4339	16 45 57.7	+43 39 27	9577 ± 118	18.3	-17.2	BCG	
152	HS 1646+4003B	16 46 35.6	+40 03 59	9877 ± 83	19.7	-15.9	BCG?	
153	HS 1649+3621 [†]	16 49 35.6	+36 21 13	9189 ± 162	18.1	-17.3	BCG	
154	HS 1650+3706	16 50 22.4	+37 06 22	6898 ± 30	19.1	-15.7	BCG	
155	HS 1653+4340	16 53 19.4	+43 40 29	59773 ± 116	+18.7	-20.9	SBN?	
156	HS 1655+3845	16 55 47.9	+38 45 56	2221 ± 112	17.1	-15.3	BCG	
157	HS 1704+3542 [‡]	17 04 51.5	+35 42 15	7602 ± 162	18.0	-17.0	BCG	
158	HS 1710+3746	17 10 19.8	+37 46 26	11196 ± 104	17.8	-18.1	BCG	
159	HS 1713+4417	17 13 12.3	+44 17 03	8425 ± 80	18.4	-16.9	BCG?	
160	HS 1713+4856	17 13 25.3	+48 56 53	8051 ± 44	19.3	-15.9	BCG	
161	HS 1716+3817	17 16 34.1	+38 17 31	13062 ± 178	16.9	-19.3	SBN?	6

^a Heliocentric velocities.; ^b absolute magnitudes are not corrected for galactic extinction;+ APM magnitudes; [†] objects from the second priority random sample; [‡] galaxies from the APM selected sample.

Table 3. (Continued)**References:**

- 1** – Klemola, A. R., Jones, B. F., & Hanson, R. B. 1987, AJ, 94, 501
- 2** – Popescu, C. C., Hopp, U., Hagen, H.-J., & Elsässer, H. 1996, A&AS, 116, 43
- 3** – Ugryumov, A. V., Pustilnik, S. A., Lipovetsky, V. A., Izotov, Yu. I., & Richter, G. M. 1998, A&AS, 131, 295
- 4** – RC3
- 5** – Weistrop, D., & Downes, R. A. 1991, AJ, 102, 1680
- 6** – Hopp, U., Engels, D., Green, R., et al. 2000, A&AS, 142, 417 (**Paper III**)
- 7** – Popescu, C. C., & Hopp, U. 2000, A&AS, 142, 247
- 8** – Falco, E. E., Kurtz, M. J., Geller, M. J., et al. 1999, PASP, 111, 438
- 9** – Markarian, B. E., Lipovetsky, V. A., Stepanian, et al. 1989, Soob. SAO. No.62, 5
- 10** – Fisher, K. B., Huchra, J. P., Strauss, M. A., et al. 1995, ApJS, 100, 69
- 11** – Grupe, D., Beuermann, K., Mannheim, K., & Thomas H.-C. 1999, A&A, 350, 895
- 12** – Kirshner, R. P., Oemler, A., Schechter, P. L., & Shectman, S. A. 1987, ApJ, 314, 493
- 13** – Tifft, W. C., Kirshner, R. P., Gregory, S. A., & Moody, J. W. 1986, ApJ, 310, 75
- 14** – Wegner, G., & Boley, F. I. 1993, AJ, 105, 660

Table 4. Emission line parameters

#	Name	$F(H\beta)^a$	$\frac{F(\lambda 3727)}{F(H\beta)}$	$\frac{F(\lambda 5007)}{F(H\beta)}$	$\frac{F(H\alpha)}{F(H\beta)}$	$\frac{F(\lambda 6583)}{F(H\alpha)}$	$\frac{F([SII])}{F(H\alpha)}$	$W_{\lambda 3727}(\text{\AA})$	$W_{H\beta}(\text{\AA})$	$W_{\lambda 5007}(\text{\AA})$
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1	HS 0809+4334	51	1.16	3.96	3.71	0.16	0.20	41	24	92
2	HS 0818+4234	+5:					0.75:		+12:	5:
3	HS 0822+4049	85	1.45	2.25	4.62	0.14	0.20	72	38	86
4	HS 0822+4238	16	2.81	3.94	4.63	0.08	0.16	39	14	55
5	HS 0824+4139	11		3.55	4.64	0.08	0.41		8	31
6	HS 0828+4305	43	1.63	1.60	5.77	0.26	0.32	16	9	15
7	HS 0828+4346	23		6.22	4.39	0.02	0.10		16	103
8	HS 0831+4208	8:		2.50	13.12	0.38	0.30		4:	11
9	HS 0849+3802 [‡]	36		2.50	7.25	0.34	0.35		7	17
10	HS 0849+3639 [‡]	20		2.10	4.45	0.08	0.26		11	22
11	HS 0850+3722 [‡]	+5:							+10:	10:
12	HS 0855+4118	64		2.59	3.88	0.12	0.25		17	45
13	HS 0855+3729 [‡]	28	1.93	0.75	4.00	0.29	0.30	24	14	11
14	HS 0900+4318	18		6.06	4.50		0.06:		19	120
15	HS 0901+3517 [‡]	2:		11:	16:		0.25:		2:	19
16	HS 0904+3752 [‡]	+13					0.08:		+26	8:
17	HS 0907+4201	4:		1.75:	9.50:	0.29	0.32		4:	7:
18	HS 0912+4427	+10				0.20	0.20:		+35	18:
19	HS 0912+4259	37	5.46	1.70	6.03	0.21	0.30	68	15	26
20	HS 0912+4433*	16		4.91	7.50	0.17	0.02:		11	53
21	HS 0912+3900 [‡]	+40				0.35	0.28		+31	2:
22	HS 0923+4021*	16		5.94	4.44	0.04	0.20		15	78
23	HS 0927+4159	30		3.13	4.23	0.09	0.29		22	75
24	HS 0934+4150	32		4.75	4.78	0.06	0.17		24	113
25	HS 0934+4256	16:		5.31:	9.25:	0.24	0.71		2:	11
26	HS 0935+4135	7:	1:	0.71:	2.86:	0.60	0.15	10:	10:	8:
27	HS 0938+4215	27		3.39	3.54	0.05	0.28		20	69
28	HS 0939+4124	61	1.43	7.89	4.39	0.01	0.04	175	171	1367
29	HS 0940+4052	133	3.65	4.48	3.73	0.05	0.17	91	41	196
30	HS 0941+4024A	12		3.11	8.09	0.40	0.40		4	13
31	HS 0941+4024B	51		2.37	4.96	0.18	0.24		17	41
32	HS 0947+3526 [‡]	24		4.29	2.92		0.20		28	118
33	HS 0948+3723*	20	1.50	5.30	2.78:	0.05:		375	32	187
34	HS 0951+3712 [‡]	6:	4.83:	2.17:	12.50:	0.32	0.36	10:	2:	5:
35	HS 0952+4024 [‡]	18		1.28	4.22	0.17	0.32		9	11
36	HS 0953+3628	77	2.53	3.83	2.64	0.10	0.19	92	42	177
37	HS 0954+3615 [‡]	6:		6.17:	6.33:	0.08	0.50:		2:	18
38	HS 0956+3702 [‡]	14	9.14	2.14	7.36	0.28	0.32	39	5	11
39	HS 0958+3654 [‡]	+95				0.53	0.44		+16	
40	HS 0958+3637*	4:		5.05:	9.77:		0.40		3:	14

Table 4. (Continued)

#	Name	$F(H\beta)^a$	$\frac{F(\lambda 3727)}{F(H\beta)}$	$\frac{F(\lambda 5007)}{F(H\beta)}$	$\frac{F(H\alpha)}{F(H\beta)}$	$\frac{F(\lambda 6583)}{F(H\alpha)}$	$\frac{F([SII])}{F(H\alpha)}$	$W_{\lambda 3727}(\text{\AA})$	$W_{H\beta}(\text{\AA})$	$W_{\lambda 5007}(\text{\AA})$
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
41	HS 0958+3718*	190	0.24	6.46	2.25	0.03		21	64	430
42	HS 1004+3730 [‡]	+81				0.42	0.40		+38	3
43	HS 1008+3620 [‡]	13		6.15	6.31	0.07	0.22		10	64
44	HS 1011+3604	26	1.81	2.08	3.85	0.16	0.29	97	15	33
45	HS 1014+3535 [‡]	5:			16:	0.37	0.25		2:	
46	HS 1016+3733*	27	3.39	4.61	5.35	0.68	0.56	23	5	22
47	HS 1021+4839	+53				0.33	0.12:		+33	6:
48	HS 1025+3707A	37		3.78	2.92	0.07	0.12		25	99
49	HS 1025+3707B	+42				0.43:			+62	
50	HS 1025+3658 [‡]	24		6.08	4.75	0.11	0.18		17	106
51	HS 1025+4057*	16:			2.19:	0.29	0.49:		19:	
52	HS 1027+4728	+24				0.42:	0.21:		+11	
53	HS 1028+3716*	32	1.00	2.88	5.63	0.20	0.30	60	13	35
54	HS 1032+4600*	45		2.88	5.81	0.15	0.32		11	34
55	HS 1043+3852 [‡]	5:		7.60:	5.60:		0.11:		6:	54
56	HS 1122+3800 [‡]	2:		9.50:	11.00:	0.05:	0.09:		6:	57
57	HS 1148+3732*	37	2.54	2.16	2.59	0.21	0.35	50	21	47
58	HS 1159+3940	14	4.07	1.79	2.71	0.21	0.18:	61	21	40
59	HS 1200+3824	+12				0.25:			+16	
60	HS 1201+3535	14	2.86	5.79	3.14	0.02:	0.20:	50	28	180
61	HS 1202+3732	27	4.66	4.60	3.50	0.04:		43	12	54
62	HS 1204+3945*	6:		4.45:	9.06:	0.10	0.24		5:	23
63	HS 1205+3940	33	1.67	5.55	3.36	0.05	0.09	81	36	237
64	HS 1206+4012	+9:							+10:	
65	HS 1210+3831*	39	4.64	2.51	3.51	0.14	0.36	59	12	32
66	HS 1211+3807*	40		2.75	5.63	0.12	0.18		8	22
67	HS 1211+4012	12	5.33	6.67	3.75	0.04:	0.13:	60	11	79
68	HS 1212+3640*	70	3.43	2.54	6.27	0.08	0.25	79	14	36
69	HS 1213+3636A	237	0.41	3.48	2.55	0.05	0.08	66	281	1094
70	HS 1213+3636B	31		3.43	3.96	0.04:	0.18		35	145
71	HS 1215+3850*	97	3.05	3.34	4.54	0.07	0.22	52	16	53
72	HS 1216+3743*	+39					0.23		+65	31
73	HS 1218+3945*	28	3.46	3.79	2.29	0.14	0.19	173	28	106
74	HS 1220+3845	19	1.27	1.43	2.85	0.62	0.21	13	13	19
75	HS 1223+3938*	48		6.25	4.46	0.07	0.15		17	109
76	HS 1224+4027	22	3.82	3.50	3.50	0.09	0.19	50	17	60
77	HS 1225+4129	17	2.47	3.06	3.82	0.14	0.38	32	14	44
78	HS 1226+4113	21	2.19	5.10	5.19	0.05:	0.12:	65	12	60
79	HS 1232+3834*	16	3.19	8.06	4.63	0.09:	0.34:	130	7	62
80	HS 1232+3947*	87		3.54	2.90		0.15		36	136
81	HS 1235+4232*	7:		2.14:	1.86:	0.15:	0.38:		24:	55:
82	HS 1236+3821*	55	5.82	5.25	5.47	0.02	0.19	124	7	41

Table 4. (Continued)

#	Name	$F(H\beta)^a$	$\frac{F(\lambda 3727)}{F(H\beta)}$	$\frac{F(\lambda 5007)}{F(H\beta)}$	$\frac{F(H\alpha)}{F(H\beta)}$	$\frac{F(\lambda 6583)}{F(H\alpha)}$	$\frac{F([SII])}{F(H\alpha)}$	$W_{\lambda 3727}(\text{\AA})$	$W_{H\beta}(\text{\AA})$	$W_{\lambda 5007}(\text{\AA})$
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
83	HS 1236+4532	+3:							+3:	
84	HS 1237+3900	40	2.33	5.35	2.90	0.07:	0.11:	79	39	227
85	HS 1238+4229	15	5.07	4.93	4.53	0.09	0.25:	86	18	91
86	HS 1242+4058A*	47		2.94	4.09	0.08	0.27		18	53
87	HS 1244+3647*	110	0.89	4.01	2.55	0.03	0.32	22	37	150
88	HS 1245+4413	32		2.16	2.06	0.18	0.26		24	57
89	HS 1300+4225	15:	2.20:	2.20:	2.33:			57	28:	64
90	HS 1302+4058	32	6.44	3.75	1.88		0.22	137	29	117
91	HS 1302+4101	22	2.09	3.59	3.45	0.09	0.20	55	25	98
92	HS 1305+4049	32	2.11	3.35	2.20	0.10	0.16	36	21	77
93	HS 1306+4146	64	1.55	5.98	2.56	0.02:	0.09:	237	146	889
94	HS 1306+4151	7:	2.71:	1.00:				17:	6:	6:
95	HS 1306+4232*	18		2.39	2.78		0.50		13	31
96	HS 1307+4225	26	0.83	5.52	3.86		0.10:	18	31	164
97	HS 1309+4446	19		2.00	3.11		0.32		9	19
98	HS 1310+4224	14	5.79	2.07	3.07		0.30	219	9	21
99	HS 1311+3628	448	1.29	5.18	2.92	0.05	0.09	299	393	2100
100	HS 1311+4507	52	2.58	4.33	2.75	0.08	0.15	116	76	335
101	HS 1311+4204	18	4.22	2.44	3.00	0.19	0.26	66	16	40
102	HS 1311+3924	179	1.89	4.38	2.30	0.04	0.13	97	61	290
103	HS 1313+4440	140		2.05	4.97	0.49	0.63		12	25
104	HS 1313+4246	30	1.93	4.80	3.57	0.05:	0.13:	57	36	180
105	HS 1313+4041*	28	2.82	5.07	3.50	0.05:	0.13:	44	19	102
106	HS 1313+4521	117	1.19	3.84	3.11	0.03:	0.12	92	77	306
107	HS 1314+4145	214		0.82	4.07	0.34	0.16		44	38
108	HS 1315+4129	+25				0.32:	0.40:	15:	+16	
109	HS 1316+3839*	48	3.13	3.17	3.29	0.13	0.29	83	25	78
110	HS 1317+4521B*	48		0.48	1.89	0.14:	0.05:		24	12
111	HS 1322+3708* [†]	20	2.60	4.35	2.85	0.16	0.19	86	35	155
112	HS 1323+3946*	103	1.80	2.32	3.79	0.31	0.15	53	15	35
113	HS 1330+3651*	188	1.92	5.02	2.71	0.01	0.13	504	84	456
114	HS 1334+3957*	60	1.28	5.22	3.12	0.02	0.10	99	96	548
115	HS 1336+3650	36		3.80	4.76	0.07	0.34		14	54
116	HS 1354+3635*	154	3.01	3.36	4.22	0.06	0.22	159	23	77
117	HS 1402+3650	107	1.38	3.03	3.27	0.12	0.21	61	27	86
118	HS 1408+4227	120	0.82	7.58	2.78	0.02:	0.04:	99	219	1615
119	HS 1408+4522*	125		2.97	2.96	0.12	0.24		35	110
120	HS 1410+3627*	13:		3.46:	4.46:	0.05:	0.26		13:	41
121	HS 1412+4203*	+16				0.44	0.25	13:	+19	
122	HS 1414+4034	8:	4:	5.13:	3.25:	0.08:	0.15:	41	12:	69
123	HS 1417+4433	44	2.18	5.73	3.07	0.02:	0.12:	167	61	364

Table 4. (Continued)

#	Name	$F(H\beta)^a$	$\frac{F(\lambda 3727)}{F(H\beta)}$	$\frac{F(\lambda 5007)}{F(H\beta)}$	$\frac{F(H\alpha)}{F(H\beta)}$	$\frac{F(\lambda 6583)}{F(H\alpha)}$	$\frac{F([SII])}{F(H\alpha)}$	$W_{\lambda 3727}(\text{\AA})$	$W_{H\beta}(\text{\AA})$	$W_{\lambda 5007}(\text{\AA})$
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
124	HS 1419+3639	69	3.74	2.43	1.96	0.13	0.31	186	41	102
125	HS 1420+3726 [†]	34	2.53	3.21	2.56		0.34	69	35	120
126	HS 1423+3945 [†]	70		4.56	2.24	0.03	0.11		136	638
127	HS 1423+4054	25	4.12	2.72	1.96	0.08	0.20	64	18	54
128	HS 1427+4309*	34	2.24	4.09	2.91	0.08	0.26	54	23	99
129	HS 1428+4056	+13				0.08:	0.15:		+21	7:
130	HS 1430+4040	121	2.94	0.79	2.78:	0.13:	0.05:	281	51	45
131	HS 1433+4411	45		3.27	3.58	0.06	0.24		10	34
132	HS 1433+4245*	+97				1.20		17:	+41	
133	HS 1437+4002	30	1.47	6.13	3.53	0.03:	0.11:	114	60	340
134	HS 1441+4122*	22	5.16	2.85	5.09	0.15	0.26	47	11	32
135	HS 1443+4027	61	1.87	2.97	3.26	0.15	0.19	163	19	56
136	HS 1447+4402	8:	3.75:	1.50:	5.50:	0.25:	0.30:	22:	5:	8:
137	HS 1448+4518*	+53				0.15	0.31	19	+44	8:
138	HS 1450+3844*	72	2.31	3.46	3.53	0.07	0.21	132	23	82
139	HS 1450+4234	29		0.72	2.93	0.27	0.40		13	10
140	HS 1458+4129	+28				0.17	0.40		+29	
141	HS 1514+4602*	+62				0.46	0.19:		+61	
142	HS 1537+4411*	19		3.91	3.30	0.03	0.23		12	50
143	HS 1545+4055	161	1.27	5.66	2.70	0.02:	0.05:	181	163	983
144	HS 1545+4459	12	7.92	3.75	5.33	0.13	0.28	76	7	29
145	HS 1546+4319*	41	3.80	1.81	4.23	0.33	0.24	25	7	14
146	HS 1603+3759 [‡]	19	2.16	4.89	2.42			104	43	228
147	HS 1613+3902	19		4.37	2.84		0.13		12	52
148	HS 1616+3627 [†]	120		0.63	3.57	0.16			11	8
149	HS 1625+3847	7:		3.00:	4.57:		0.25		8:	25
150	HS 1633+4703*	170	3.96	3.29	3.38	0.07	0.23	76	16	55
151	HS 1645+4339	23		4.43	4.83	0.06	0.22		11	51
152	HS 1646+4003B	21:		1.62:	3.33:	0.14			13:	19:
153	HS 1649+3621 [†]	21:	1.86:	3.57:	3.43:			37:	13:	48
154	HS 1650+3706	66	2.36	5.55	2.39	0.08	0.09	275	124	647
155	HS 1653+4340	6:	3.83:	3.17:	6.00:	0.44:	0.44:	21:	4:	13:
156	HS 1655+3845*	49	2.31	4.14	2.76		0.19	60	30	125
157	HS 1704+3542 [‡]	4:	12.54:	8.16:	6.16:		0.22:	60	5:	46
158	HS 1710+3746*	19	2.84	3.11	3.74	0.07	0.23	110	24	68
159	HS 1713+4417	15	4.27	3.07	5.93	0.22	0.25	24	5	16
160	HS 1713+4856	140	1.04	5.91	3.08	0.04	0.05	147	101	632
161	HS 1716+3817	9:		1.67:	10.11:	0.13	0.32		3:	5:

^a Flux in 10^{-16} ergs s⁻¹ cm⁻²; * objects observed during non-photometric conditions;[†] objects from the second priority random sample; [‡] galaxies from the APM selected sample;⁺ parameters for H α emission line; : parameters with less confident values.

Table 5. Coordinates, Redshifts and Magnitudes of QSOs

#	Name	α (1950)	δ (1950)	z^a	m_B	Detected emission lines
	(1)	(2)	(3)	(4)	(5)	(6)
1	HS 0936+4204	09 36 59.6	+42 04 24	3.3211	18.3	Ly α 1216 Å, Siiv/Oiv] 1400 Å, Civ 1549 Å
2	HS 1009+3706	10 09 02.4	+37 06 31	3.0393	17.6	Ly α 1216 Å, Siiv/Oiv] 1400 Å, Civ 1549 Å
3	HS 1224+4410	12 24 55.1	+44 10 03	3.0718	19.5	Ly α 1216 Å, Civ 1549 Å
4	HS 1301+4233	13 01 47.0	+42 33 30	0.8460:	+18.7	MgII 2798 Å
5	HS 1541+4452	15 41 23.0	+44 52 27	2.3588	18.8	Ly α 1216 Å, Siiv/Oiv] 1400 Å, Civ 1549 Å

^a Observed redshift; + APM magnitudes; : less confident values.

Table 6. Galaxies without detected emission lines

#	Name	α (1950)	δ (1950)	v_0^a	m_B	M_B^b	Absorption lines
	(1)	(2)	(3)	(4)	(5)	(6)	(7)
1	HS 1008+3538	10 08 00.1	+35 38 02	13968:	17.4	-19.0	H β , MgIb, H α
2	HS 1207+3745	12 07 38.0	+37 45 53	13381:	18.8	-17.5	H β , H α
3	HS 1408+3524 [†]	14 08 32.6	+35 24 23	19933	16.8	-20.3	CaK, CaH, G _{band} , H β , MgIb, CaFe5269, NaD
4	HS 1419+4026	14 19 50.0	+40 26 28	14235:	18.1	-18.3	G _{band} , H α

^a Heliocentric velocities; ^b absolute magnitudes are not corrected for galactic extinction;

[†] objects from the second priority random sample; : less confident values.

Table 7. Stars

#	Name	α (1950)	δ (1950)	m_B	Type	Spectral features
	(1)	(2)	(3)	(4)	(5)	(6)
1	HS 0852+4257	08 52 23.3	+42 57 22	+20.2	A	H_γ , H_β , H_α
2	HS 0945+4851	09 45 00.5	+48 51 00	+18.0	F-G	H_β , MgIb, CaFe5269, H_α
3	HS 1012+3815	10 12 45.4	+38 15 01	17.2	A	H_γ , H_β , H_α
4	HS 1013+3804 [†]	10 13 05.6	+38 04 27	17.7	F	H_β , NaD, H_α
5	HS 1029+4020 [†]	10 29 10.6	+40 20 35	16.7	G	MgIb, CaFe5269, NaD, H_α
6	HS 1158+3920	11 58 04.6	+39 20 32	+18.2	A-F	H_γ , H_β , MgIb, NaD, H_α
7	HS 1411+4312	14 11 12.8	+43 12 53	19.6	G	G_{band} , MgIb, NaD, H_α
8	HS 1424+4029	14 24 57.9	+40 29 28	19.8	F	G_{band} , H_β , MgIb, NaD, H_α
9	HS 1609+4941	16 09 46.6	+49 41 45	19.1	G	G_{band} , MgIb, CaFe5269, H_α
10	HS 1630+3620	16 30 33.0	+36 20 17	19.1	G	G_{band} , MgIb, CaFe5269, NaD, H_α
11	HS 1640+4158	16 40 18.0	+41 58 42	18.3	G	G_{band} , MgIb, NaD
12	HS 1652+3820	16 52 47.9	+38 20 16	17.7	G	G_{band} , MgIb, CaFe5269, NaD, H_α
13	HS 1653+4430	16 53 49.9	+44 30 10	19.7	F	CaK, CaH, G_{band} , H_β , MgIb, NaD, H_α
14	HS 1656+4309	16 56 02.7	+43 09 01	19.2	A	H_γ , H_β , H_α
15	HS 1657+4419	16 57 28.6	+44 19 26	20.0	F	CaK, CaH, H_γ , H_β , H_α
16	HS 1710+4343	17 10 04.2	+43 43 47	+18.2	F	H_β , MgIb, H_α
17	HS 1711+3804	17 11 31.4	+38 04 32	+19.1	F	H_β , MgIb, NaD, H_α
18	HS 1716+3544	17 16 14.0	+35 44 07	18.9	F	G_{band} , H_γ , H_β , MgIb, CaFe5269, H_α
19	HS 1720+3927	17 20 13.6	+39 27 11	18.1	F-G	G_{band} , H_β , MgIb, NaD, H_α
20	HS 1733+3749	17 33 41.9	+37 49 13	18.4	G	G_{band} , H_β , MgIb, NaD, H_α
21	HS 1733+3915 [‡]	17 33 45.8	+39 15 08	18.1	F-G	H_β , MgIb, NaD, H_α

[†] objects from the second priority random sample; [‡] galaxies from the APM selected sample;

⁺ APM magnitudes.

This figure "A01.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A02.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A03.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A04.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A05.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A06.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A07.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A08.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A09.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A10.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A11.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A12.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A13.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A14.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A15.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A16.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>

This figure "A17.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0106186v1>